



**"Gheorghe Asachi" Technical University of Iasi, Romania**



---

## **SPATIAL DISTRIBUTION AND SOURCE APPORTIONMENT OF POTENTIALLY TOXIC ELEMENTS IN DILUVIAL AND ALLUVIAL TOPSOIL AROUND XIAOLONGTAN LIGNITE DEPOSIT, GEJIU YUNNAN PROVINCE, CHINA**

**Shuran Yang<sup>1,2\*</sup>, Zailing Cheng<sup>1,2</sup>, Qianrui Huang<sup>1,2</sup>, Xianfeng Cheng<sup>1,2</sup>, Yu Cheng<sup>3,4</sup>**

<sup>1</sup>Yunnan Land and Resources Vocational College, Kunming 652501, Yangzonghai Scenic Area, Yunnan Province, China

<sup>2</sup>Engineering Center of Yunnan Education Department for Health Geological Survey & Evaluation

<sup>3</sup>International Research Center of Big Data for Sustainable Development Goals, Beijing 100094, China

<sup>4</sup>Key Laboratory of Digital Earth Science, Aerospace Information Research Institute, Chinese Academy of Sciences, Beijing 100094, China

---

### **Abstract**

Lignite mining can significantly increase potentially toxic elements (PTEs) in soils through excavation and coal combustion. This study measured the concentrations of As, Hg, Cd, Cu, Pb, and Zn in diluvial and alluvial topsoil, as well as lignite samples, from the Xiaolongtan Lignite Deposit (XLD) in Gejiu, Yunnan Province, China. GIS mapping and absolute principle component scores -multivariate linear regression (APCS-MLR) model were employed to determine the spatial distribution, identify pollution sources, and assess source-specific ecological and health risks. Topsoil PTEs concentrations (mg/kg) ranged from 5.78–117 (As), 0.03–0.30 (Hg), 0.12–2.07 (Cd), 6.10–5221 (Cu), 3.28–978 (Pb), and 15.1–2620 (Zn), exceeding local background levels. Lignite contained (mg/kg) 16.8 (As), 0.07 (Hg), 0.72 (Cd), 44.2 (Cu), 58.2 (Pb), and 134 (Zn), higher than national and global averages. Enrichment factors (EF) followed the order Cu > As > Cd > Zn > Pb, suggesting anthropogenic inputs. Two principal components were identified: PC1 (68.26%) associated with Cu, Pb, Zn, Cd from agriculture and traffic; and PC2 (31.74%) associated with As and Hg from mining, industry, and weathering. The total ecological risk was extremely high (RI = 5817.15), with PC2 contributing more (25.45%) than PC1 (22.74%). Non-carcinogenic risks were significant, with total hazard index (THI) values of 43.8 (children) and 19.6 (adults); carcinogenic risks were also high (TCR =  $5.1 \times 10^{-3}$  for children and  $2.2 \times 10^{-3}$  for adults). Overall, mining and industrial activities were the major sources of ecological and health risks, with As posing the greatest threat, requiring enhanced management and control.

**Key words:** APCS-MLR model, Potential toxic elements, source apportionment, Source-specific risk assessment, Xiaolongtan Lignite Deposit

*Received: April, 2024; Revised final: June, 2025; Accepted: June, 2025*

---

---

\* Author to whom all correspondence should be addressed: e-mail: yangshuran1988@foxmail.com, Phone: +8618669001620, Fax: +087168160123