



“Gheorghe Asachi” Technical University of Iasi, Romania



ANALYSIS OF THE CONTINUOUS MEASUREMENTS OF PM₁₀ AND PM_{2.5} CONCENTRATIONS IN BEIRUT, LEBANON

Wehbeh Farah^{1*}, Myriam Mrad Nakhle^{2,5}, Maher Abboud³, Nelly Ziade⁴, Isabella Annesi-Maesano⁵, Rita Zaarour⁶, Nada Saliba⁶, Georges Germanos¹, Najat Aoun Saliba⁷, Alan L. Shihadeh⁸, Jocelyne Gerard⁶

¹Physics Department, Saint Joseph University of Beirut, Beirut, Lebanon

²Biology Department, Saint Joseph University of Beirut, Beirut, Lebanon

³Chemistry Department, Saint Joseph University of Beirut, Beirut, Lebanon

⁴Rheumatology Department, Saint Joseph University of Beirut, Beirut, Lebanon

⁵EPAR, Institute Pierre Louis of Epidemiology and Public Health, UMR-S 1136 INSERM & UPMC Paris 6, Sorbonnes Universities, Medical School Saint-Antoine, Paris, France

⁶Geography Department, Saint Joseph University of Beirut, Beirut, Lebanon

⁷Mechanical Engineering Department, American University of Beirut, Beirut, Lebanon

⁸Chemistry Department, American University of Beirut, Beirut, Lebanon

Abstract

Atmospheric concentrations of PM_{2.5} and PM₁₀ were measured in Beirut, Lebanon, for a period of 12 months. The daily average concentrations of PM₁₀ and PM_{2.5} were found to be 51.3 ± 33.1 and $30.3 \pm 9.4 \mu\text{g}\cdot\text{m}^{-3}$, respectively, with corresponding maximum values of 359.7 and 208.6 $\mu\text{g}\cdot\text{m}^{-3}$. The annual average concentrations of PM₁₀ and PM_{2.5} exceeded the World Health Organization's standards by 150% and 200%, respectively. The mean concentration of coarse particles (PM_{10-2.5}) was found to be 41% of the average PM₁₀, suggesting that the site was also influenced by re-suspended surface dust and soil. The mean PM_{2.5}/PM₁₀ ratio for the entire study period was 0.61 ± 0.12 . This indicates that in Beirut, PM_{2.5} accounts for about 61% of PM₁₀. Such a large fraction of fine particles could have considerable effect on health; thus, it is necessary to quantify its impact. Daily concentrations of PM₁₀ and PM_{2.5} exceeded the upper threshold limit on 133 and 129 days, respectively, representing 39% and 38% of the entire sample, respectively. These findings indicate the important role dust events play within this area. Concentrations of PM_{2.5} were highly correlated with NO₂, whereas concentrations of PM₁₀ and PM_{10-2.5} were not associated with any gaseous pollutant. Regression analysis showed that 93% of PM_{2.5} and 43% of PM₁₀ particle mass concentrations were derived from road traffic exhaust in Beirut.

Key words: air quality, dust event, health effect, particulate matter, regression analysis

Received: October, 2013; *Revised final:* September, 2014; *Accepted:* September, 2014

* Author to whom all correspondence should be addressed: e-mail: wehbeh.farah@usj.edu.lb; Phone: +96 1142 1374; Fax: +96 1453 2657