



“Gheorghe Asachi” Technical University of Iasi, Romania



---

## COMPOSITION OF TRACE METALS IN INDOOR DUST DURING AND AFTER BUILDING RENOVATION

Mohd Talib Latif<sup>1\*</sup>, Tan Sheh Na<sup>1</sup>, Nor Hafizah Baharudin<sup>1</sup>, Thunwadee Srithawirat<sup>2</sup>,  
Noorlin Mohamad<sup>3</sup>, Md Firoz Khan<sup>4</sup>, Fazrul Razman Sulaiman<sup>5</sup>

<sup>1</sup>School of Environmental and Natural Resource Sciences, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, 43000, Bangi, Selangor, Malaysia

<sup>2</sup>Department of Environmental Science, Faculty of Science and Technology, Pibulsongkram Rajabhat University, Phitsanulok 65000, Thailand

<sup>3</sup>Environmental Research Group, School of Marine and Environmental Sciences, Universiti Malaysia Terengganu, 21030 Kuala Terengganu, Terengganu, Malaysia

<sup>4</sup>Centre for Tropical Climate Change System (IKLIM), Institute for Climate Change, Universiti Kebangsaan Malaysia, 43600 Bangi, Selangor, Malaysia

<sup>5</sup>Faculty of Applied Sciences, Universiti Teknologi MARA Cawangan Pahang, 26400 Bandar Tun Abdul Razak Jengka, Pahang, Malaysia

---

### Abstract

A study was conducted to determine the composition and concentration of trace metals in suspended particulate matter (PM<sub>10</sub>) and indoor dust in two academic buildings of Universiti Kebangsaan Malaysia. One building was in the final phases of renovation (UR) and the other building had just finished renovation (AR). PM<sub>10</sub> sampling of air was performed using a low volume sampler (LVS), and samples of indoor dust were obtained using a brush and a small plastic bag. The compositions of trace metals in both samples were determined by inductively coupled plasma mass spectrometry (ICP-MS). PM<sub>10</sub> concentrations in the building under renovation (UR) were found to be higher ( $77.1 \pm 32.4 \mu\text{g m}^{-3}$  to  $355.4 \pm 38.3 \mu\text{g m}^{-3}$ ) than those in the building after renovation (AR) ( $91.7 \pm 41.2 \mu\text{g m}^{-3}$  to  $147.9 \pm 3.0 \mu\text{g m}^{-3}$ ). Higher concentrations of PM<sub>10</sub> were found in the open areas (foyer and corridor) than in closed areas (classroom and laboratory). The concentrations of trace metals in PM<sub>10</sub> and indoor dust were found to be dominated by Zn, followed by Pb > Cu > Cd. There is no indication of health adverse effect based on the concentration of trace metals recorded in the UR and AR buildings from this study.

*Key words:* health risk, indoor dust, PM<sub>10</sub>, renovation, trace metals

*Received:* June, 2013; *Revised final:* October, 2014; *Accepted:* October, 2014; *Published in final edited form:* August, 2018

---

\* Author to whom all correspondence should be addressed: e-mail: talib@ukm.edu.my; Phone: +60389215433; Fax: +60389255104