Environmental Engineering and Management Journal

March 2019, Vol. 18, No. 3, 555-562 http://www.eemj.icpm.tuiasi.ro/; http://www.eemj.eu



"Gheorghe Asachi" Technical University of Iasi, Romania



SPATIAL EVALUATION OF NO₂, SO₂, PM10 AND BTEX CONCENTRATIONS IN URBAN AMBIENT AIR

Dainius Paliulis*, Pranas Baltrėnas

Institute of Environmental Protection, Vilnius Gediminas Technical University, Saulėtekio al. 11, Vilnius, LT–10223, Lithuania

Abstract

During the 2014-2015 period of environmental air quality research, concentrations of NO₂, SO₂ and BTEX (benzene, toluene, ethylbenzene and xylenes) and PM10 were measured. In order to evaluate the spatial distribution of pollutants, 5 sampling sites in Anykščiai city (Lithuania) were chosen. Measured values of NO₂, SO₂, PM10 and benzene were compared with limit values set for these pollutants in ambient air according to 2008/50/EC and 2000/69/EC directives. Toluene, ethylbenzene, and xylenes (*ortho, meta*, and *para*) were not included into the list of pollutants, which amount in ambient air is regulated according to European Union criteria. These pollutants are regulated since July 1, 2007 according to national criteria of Lithuania. The NO₂ concentration was found in the range of 10.0–29.0 μ g/m³. Benzene concentrations in Anykščiai varied from 0.4–2.8 μ g/m³, ethylbenzene concentrations in Anykščiai varied from 0.9–2.8 μ g/m³. The annual average of NO₂, SO₂, PM10 and BTEX and concentrations were below the permissible limits in all five places.

Key words: ambient air, BTEX, NO2, SO2, PM10

Received: December, 2016; Revised final: February, 2017; Accepted: March, 2017; Published in final edited form: March, 2019

^{*}Author to whom all correspondence should be addressed: e-mail: dainius.paliulis@vgtu.lt; Phone +370 5 2744725; Fax. +370 5 2744726