



“Gheorghe Asachi” Technical University of Iasi, Romania



UV-H₂O₂ ADVANCED OXIDATION OF ANIONIC SURFACTANT: REACTION KINETICS, EFFECTS OF INTERFERING SUBSTANCES AND OPERATING CONDITIONS

Bijoli Mondal¹, Asok Adak^{1*}, Pallab Datta²

¹Department of Civil Engineering, Indian Institute of Engineering Science and Technology,
Shibpur, Howrah 711103, India

²Center for Healthcare Science and Technology, Indian Institute of Engineering Science and Technology,
Shibpur, Howrah 711103, India

Abstract

This paper deals with the degradation kinetics of an anionic surfactant (sodium dodecyl sulfate, SDS) under UV and UV-H₂O₂ advanced oxidation process (AOP). Experiments were performed in a batch reactor which emits monochromatic light centered at 253.7 nm. The photon flux of the UV reactor was $1.9(\pm 0.1) \times 10^{-4}$ Einstein/L-min. Under direct UV, only 45% SDS degradation was observed at a fluence of 40.65 J/cm² for initial SDS concentration of 0.35 mM. The apparent fluence-based pseudo-first order rate constant was found to be $1.77(\pm 0.24) \times 10^{-5}$ cm²/mJ. However, very fast degradation (almost 100%) was observed for UV-H₂O₂ process at a fluence of 0.45 J/cm². The apparent fluence-based rate constant in this case was found to be about 450 times higher than direct photolysis. Further, using a probe compound (para chlorobenzoic acid) competition kinetics study was performed to determine the second order hydroxyl radical rate constant for SDS. The rate constant thus obtained was $8.18 (\pm 0.26) \times 10^9$ M⁻¹ s⁻¹. Effects of different operating parameters like H₂O₂ concentration, initial SDS concentration and water quality parameters, including pH (7 - 12), alkalinity (0 - 10 mM HCO₃⁻) and nitrate concentration (0 - 1 mM as NO₃⁻), were investigated. The observed rate constants were influenced by all the studied parameters and these parameters were optimized for field application. Degradation of SDS was also carried out in municipal wastewater. In conclusion, UV-H₂O₂ AOP is an efficient treatment method for SDS in environmental matrices.

Key words: direct photolysis, hydroxyl radical rate constant, scavenging effect, sodium dodecyl sulfate, wastewater

Received: May, 2017; *Revised final:* November, 2017; *Accepted:* January, 2018; *Published in final edited form:* June 2019

* Author to whom all correspondence should be addressed: e-mail: asok@civil.iests.ac.in; Phone: +91-33-2668 4561 (extn. 713);
Fax: +91-33-2668 2916