



“Gheorghe Asachi” Technical University of Iasi, Romania



## VARIATION IN COMPOST PEDOFAUNA UNDER DIFFERENT CARBON INPUTS AND WATERING TREATMENTS

Fatiha Metna<sup>1</sup>, Arezki Hammoum<sup>1</sup>, Rachid Slimani<sup>1,2\*</sup>, Melissa Kanane<sup>1</sup>

<sup>1</sup>Laboratory of Natural Resources, Faculty of Biological and agricultural Sciences,  
Mouloud Mammeri University of Tizi-Ouzou, Algeria

<sup>2</sup>Department of Biology, Faculty of Sciences, M'Hamed Bougara University of Boumerdes, Algeria

### Abstract

The current study was carried out at Bouzeguène's psychoeducational center (Tizi Ouzou, Algeria), with a focus on the monitoring of compost pedofauna in relation to carbon inputs (dead leaves, paper-cardboard, and olive pomace) and watering liquids (water, lactoserum, and vegetation water). These composts were created by valorizing the organic fraction, which constituted 60 to 70% of household and similar waste. Nine tests were run to monitor the composting process. Two tons of organic waste were combined with 1.5 tons of three kinds of carbon waste. This mixture was composed of 50% nitrogenous matter and 50% carbon material. Three composters were monitored for each type of carbon material: the first was watered with water, the second with vegetation water, and the third with lactoserum. The analyzed composts contained eleven faunal groups. The average number of individuals varied between 356 ind/6250 cm<sup>3</sup> of compost mixed with dead leaves and watered with vegetation water and 9286 ind/6250 cm<sup>3</sup> of compost mixed with cardboard paper and watered with lactoserum. Mites dominated the nine substrates, with an average density of 99% in composts containing cardboard paper, as well as those mixed with olive pomace, and watered with lactoserum. They were followed by *Diptera* and *Coleoptera* larvae, with the remaining faunal groups under represented or absent, most likely due to inter and intraspecific competition and predation. Our findings revealed that watering with vegetation water encouraged the formation of a larger number of faunal groups.

**Key words:** Bouzeguène, composting, organic waste, pedofauna

Received: March, 2023; Revised final: January, 2025; Accepted: January, 2025; Published in final edited form: September, 2025

\* Author to whom all correspondence should be addressed: e-mail: r.slimani@univ-boumerdes.dz; Phone: +213 0664 08 81 39