



ASSESSMENT OF DEUTERIUM DEPLETION IN PLYMOUTH ROCK CHICKENS' BODIES

Ioan Ștefănescu¹, Nicolae Manolescu², Gheorghe Tițescu¹, Emilia Balint³, Radu Tamaian^{1*}, Victor Feurdean⁴

¹*National Research and Development Institute for Cryogenics and Isotopic Technologies, 4 Uzinei Street,
240050 Râmnicu Vâlcea, Romania*

²*Oncology Institute "Prof. Dr. Al. Trestioreanu", 252 Fundeni Av., 022338 Bucharest, Romania*

³*Comparative Medicine Institute, 105 Splaiul Independenței Av., 0500907 Bucharest, Romania*

⁴*National Institute for Research and Development of Isotopic and Molecular Technologies, 65-103 Donath Street,
400293 Cluj-Napoca, Romania*

Abstract

Deuterium content of living animals body can be increased, with dramatical effect over the health state, through bioaccumulation process by continuously adding of heavy water in their nourishment. On the other hand, deuterium depletion has been mentioned in literature as a health stimulator and a new possible anticancer tool. In this context, the present paper investigated the deuterium depletion process at chickens (breed Plymouth Rock, white variety), after DDW's (deuterium depleted water) administration. Chickens were raised and nourished with standard food and drinking water, respectively DDW. Liquid and solid samples were analyzed through mass-spectrometry for isotopic deuterium content. The results showed that DDW's administration decreased the isotopic content in deuterium in chickens' bodies. The deuterium depletion is made in time and it is a process which affects the entire organism.

Key words: chickens, depletion, deuterium

Received: September, 2010; Revised: November, 2010; Accepted: November, 2010

* Author to whom all correspondence should be addressed: e-mail: tradu@icsi.ro; Phone +40250 732744, Fax +40250 732746